

Detection of some elements in the stable homotopy groups of spheres *

LIU XIUGUI

School of Mathematical Sciences and LPMC

Nankai University

Tianjin 300071, P. R. China

E-mail: xgliu@nankai.edu.cn

Abstract In this paper we constructs a new nontrivial family in the stable homotopy groups of spheres $\pi_{p^n q + 2pq + q - 3} S$ which is of order p and is represented by $k_0 h_n \in Ext_A^{3, p^n q + 2pq + q}(\mathbb{Z}_p, \mathbb{Z}_p)$ in the Adams spectral sequence, where $p \geq 5$ is an odd prime, $n \geq 3$ and $q = 2(p - 1)$. In the course of the proof, a new family of homotopy elements in $\pi_* V(1)$ which is represented by $\beta_* i'_* i_*(h_n) \in Ext_A^{2, p^n q + (p+1)q + 1}(H^* V(1), \mathbb{Z}_p)$ in the Adams sequence is detected.

Keywords Stable homotopy groups of spheres, Adams spectral sequence, May spectral sequence, Steenrod algebra.

2000 MR Subject Classification 55Q45

1 Introduction and the main results

Let A be the mod p Steenrod algebra and S be the sphere spectrum localized at an odd prime p . To determine the stable homotopy groups of spheres $\pi_* S$ is one of the central problems in homotopy theory. One of the main tools to reach it is the Adams spectral sequence:

$$E_2^{s,t} = Ext_A^{s,t}(\mathbb{Z}_p, \mathbb{Z}_p) \Rightarrow \pi_{t-s} S,$$

where the $E_2^{s,t}$ -term is the cohomology of A . So far, not so many families of homotopy elements in $\pi_* S$ have been detected. For example, a family $\varsigma_{n-1} \in \pi_{p^n q + q - 3} S$ for $n \geq 2$ which has filtration 3 in the Adams spectral sequence and is represented by

*This work was supported partially by the National Natural Science Foundation of China (Grant No. 10501045), and the Fund of the Personnel Division of Nankai University.

$h_0 b_{n-1} \in Ext_A^{3,p^n q+q}(\mathbb{Z}_p, \mathbb{Z}_p)$ has been detected in reference [1], where $q = 2(p-1)$. In this paper, we also detect a family of homotopy elements in $\pi_{p^n q+pq-3} S$ which has filtration 3 and is represented by $k_0 h_n \in Ext_A^{3,p^n q+2pq+q}(\mathbb{Z}_p, \mathbb{Z}_p)$ in the Adams spectral sequence.

From reference [2], $Ext_A^{1,*}(\mathbb{Z}_p, \mathbb{Z}_p)$ has $\mathbb{Z}_p$ -bases consisting of $a_0 \in Ext_A^{1,1}(\mathbb{Z}_p, \mathbb{Z}_p)$, $h_i \in Ext_A^{1,p^i q}(\mathbb{Z}_p, \mathbb{Z}_p)$ for all $i \geq 0$ and $Ext_A^{2,*}(\mathbb{Z}_p, \mathbb{Z}_p)$ has $\mathbb{Z}_p$ -bases consisting of α_2 , a_0^2 , $a_0 h_i (i > 0)$, $g_i (i \geq 0)$, $k_i (i \geq 0)$, $e_i (i \geq 0)$, and $h_i h_j (j \geq i+2, i \geq 0)$ whose internal degrees are $2q+1$, 2 , $p^i q+1$, $p^{i+1} q+2p^i q$, $2p^{i+1}+p^i q$, $p^{i+1} q$ and $p^i q+p^j q$ respectively.

Let M be the Moore spectrum modulo a prime $p \geq 5$ given by the cofibration

$$S \xrightarrow{p} S \xrightarrow{i} M \xrightarrow{j} \Sigma S. \quad (1.1)$$

Let $\alpha : \Sigma^q M \rightarrow M$ be the Adams map and K be its cofibre given by the cofibration

$$\Sigma^q M \xrightarrow{\alpha} M \xrightarrow{i'} K \xrightarrow{j'} \Sigma^{q+1} M, \quad (1.2)$$

where $q = 2(p-1)$. This spectrum which we briefly write as K is known to be the Toda-Smith spectrum $V(1)$. Let $V(2)$ be the cofibre of $\beta : \Sigma^{(p+1)q} K \rightarrow K$ given by the cofibration

$$\Sigma^{(p+1)q} K \xrightarrow{\beta} K \xrightarrow{\bar{i}} V(2) \xrightarrow{\bar{j}} \Sigma^{(p+1)q+1} K. \quad (1.3)$$

Our results can be stated as follows.

Theorem I *Let $p \geq 5, n \geq 3$, then*

$$\beta_* i'_* i_*(h_n) \neq 0 \in Ext_A^{2,p^n q+(p+1)q+1}(H^* K, \mathbb{Z}_p)$$

is a permanent cycle in the Adams spectral sequence and converges to a nontrivial element $\zeta_n \in \pi_{p^n q+(p+1)q-1} K$, where $h_n \in Ext_A^{1,p^n q}(\mathbb{Z}_p, \mathbb{Z}_p)$.

Theorem II *Let $p \geq 5, n \geq 3$, then*

$$k_0 h_n \neq 0 \in Ext_A^{3,p^n q+2pq+q}(\mathbb{Z}_p, \mathbb{Z}_p)$$

is a permanent cycle in the Adams spectral sequence and it converges to a nontrivial element of order p in $\pi_{p^n q+2pq+q-3} S$.

Remark: The $k_0 h_n$ -element obtained in Theorem II is an indecomposable element in $\pi_* S$, i.e., it is not a composition of elements of lower filtration in $\pi_* S$, because $h_n (n > 0)$ is known to die in the Adams spectral sequence.

After giving some useful propositions in Section 2, the proofs of the main theorems will be given in Section 3.

2 Some preliminaries on low-dimensional Ext groups

In this section, we will prove some results on Ext groups of lower dimension which will be used in the proofs of the theorems.

Proposition 2.1 *Let $p \geq 5, n \geq 3, a_0 \in Ext_A^{1,1}(\mathbb{Z}_p, \mathbb{Z}_p), h_n \in Ext_A^{1,p^ng}(\mathbb{Z}_p, \mathbb{Z}_p), b_n \in Ext_A^{2,p^{n+1}g}(\mathbb{Z}_p, \mathbb{Z}_p)$ respectively. Then we have the following:*

- (1) $Ext_A^{4,p^ng+pq+2}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{a_0a_0h_1h_n\}.$
- (2) $Ext_A^{5,p^ng+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p) = 0.$

Proof. (1) See [3, Theorem 4.1].

(2) The proof is similar to that given in the proof of [4, Proposition 1.2]. We can show that in the May spectral sequence $E_1^{5,p^ng+(p+2)q,*} = 0$. Then

$$Ext_A^{5,p^ng+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p) = 0.$$

Here the proof is omitted. □

The following lemma is used in the proofs of many propositions in this section.

First recall spectra $V(k) = \{V(k)_n\}$ for $n \geq -1$ which are so-called Toda-Smith spectra. The spectrum $V(n)$ given in [5] such that the $\mathbb{Z}_p$ -cohomology

$$H^*(V(n), \mathbb{Z}_p) \cong E(n) = E(\mathcal{Q}_0, \mathcal{Q}_1, \dots, \mathcal{Q}_n),$$

the exterior algebra generator by Milnor basis elements $\mathcal{Q}_0, \mathcal{Q}_1, \dots, \mathcal{Q}_n$ in A . The spectra $V(n)$ for $n \geq -1$ are defined inductively by $V(-1) = S$ and the cofibration

$$\Sigma^{2(p^n-1)}V(n-1) \xrightarrow{\alpha^{(n)}} V(n-1) \xrightarrow{i_n} V(n) \xrightarrow{j_n} \Sigma^{2p^n-1}V(n-1). \quad (2.1)$$

When $n = 0, 1, 2$, the above cofibration sequence just is the cofibration sequences (1.1), (1.2) and (1.3) respectively. $\alpha^{(n)}$ stand for the maps p, α, β in (1.1), (1.2) and (1.3) respectively. Here $V(-1) = S, V(0) = M, V(1) = K, i_0 = i, i_1 = i', i_2 = \bar{i}, j_0 = j, j_1 = j', j_2 = \bar{j}$. The existence of $V(n)$ is assured [5, Theorem 1.1] for $n = 1, p \geq 3$ and for $n = 2, p \geq 5$.

By the definition of Ext groups, from (2.1) we can easily have the following lemma.

Lemma 2.1 *With notations as above. We have the following two long exact sequence:*

$$\begin{aligned} (1) \quad & \dots \rightarrow Ext_A^{s-1,t-(2p^n-1)}(H^*V(n-1), \square) \xrightarrow{\alpha_*^{(n)}} Ext_A^{s,t}(H^*V(n-1), \square) \xrightarrow{(i_n)^*} \\ & Ext_A^{s,t}(H^*V(n), \square) \xrightarrow{(j_n)^*} Ext_A^{s,t-(2p^n-1)}(H^*V(n), \square) \rightarrow \dots \\ (2) \quad & \dots \rightarrow Ext_A^{s-1,t-(2p^n-1)}(\square, H^*V(n-1)) \xrightarrow{\alpha^{(n)*}} Ext_A^{s,t}(\square, H^*V(n-1)) \xrightarrow{(j_n)^*} \\ & Ext_A^{s,t}(\square, H^*V(n)) \xrightarrow{(i_n)^*} Ext_A^{s,t}(\square, H^*V(n)) \rightarrow \dots \end{aligned}$$

Here $\square$ are an arbitrary A -module. □

Proposition 2.2 *Let $p \geq 5, n \geq 3$. Then $Ext_A^{3,p^ng+(p+2)q+1}(H^*M, H^*M)$ has a unique generator $\overline{h_n g_0}$, where $\overline{h_n g_0}$ satisfies $i^*j_*\overline{h_n g_0} = h_n g_0$, the generator of $Ext_A^{3,p^ng+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p)$ stated in [6, Table 8.1].*

Proof. First consider the exact sequence

$$\begin{aligned} Ext_A^{3,p^n q+(p+2)q+1}(\mathbb{Z}_p, \mathbb{Z}_p) &\xrightarrow{j^*} Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, H^*M) \xrightarrow{i^*} \\ Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p) &\xrightarrow{p^*} Ext_A^{4,p^n q+(p+2)q+1}(\mathbb{Z}_p, \mathbb{Z}_p) \end{aligned}$$

induced by (1.1). Since we know that $Ext_A^{3,p^n q+(p+2)q+1}(\mathbb{Z}_p, \mathbb{Z}_p)$ is zero (cf. [6, Table 8.1]) and $Ext_A^{4,p^n q+(p+2)q+1}(\mathbb{Z}_p, \mathbb{Z}_p)$ is zero (cf. [7, Proposition 2.1]), the above i^* is an isomorphism. Then we see that $Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, H^*M)$ has unique generator $\overline{h_n g_0}$, where $\overline{h_n g_0}$ satisfies $i^* \overline{h_n g_0} = h_n g_0$, the unique generator of $Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p)$ stated in [6, Table 8.1].

At last, look at the following exact sequence induced by (1.1)

$$\begin{aligned} Ext_A^{3,p^n q+(p+2)q+1}(\mathbb{Z}_p, H^*M) &\xrightarrow{i_*} Ext_A^{3,p^n q+(p+2)q+1}(H^*M, H^*M) \xrightarrow{j_*} \\ Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, H^*M) &\xrightarrow{p_*} Ext_A^{4,p^n q+(p+2)q+1}(\mathbb{Z}_p, H^*M). \end{aligned}$$

Since the first group is zero by virtue of $Ext_A^{3,p^n q+(p+2)q+r}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $r = 1, 2$ (cf. [6, Table 8.1]) and the forth group is zero by virtue of the facts that $Ext_A^{4,p^n q+(p+2)q+t}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $t = 1, 2$ (cf. [7, Proposition 2.1]), then the above j_* is an isomorphism. Thus $Ext_A^{3,p^n q+(p+2)q+1}(H^*M, H^*M)$ has a unique generator $\overline{\overline{h_n g_0}}$, where $\overline{\overline{h_n g_0}}$ satisfies $j_* \overline{\overline{h_n g_0}} = \overline{h_n g_0}$. This finishes the proof of Proposition 2. $\square$

Proposition 2.3 *Let $p \geq 5, n \geq 3$, then*

$$Ext_A^{3,p^n q+(p+2)q}(H^*M, H^*M) \cong \mathbb{Z}_p \{i_* j_* \overline{\overline{h_n g_0}}, j^* i^* \overline{\overline{h_n g_0}}\}.$$

Proof. Consider the exact sequence

$$\begin{aligned} Ext_A^{2,p^n q+(p+2)q-1}(\mathbb{Z}_p, \mathbb{Z}_p) &\xrightarrow{p^*} Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p) \xrightarrow{j^*} \\ Ext_A^{3,p^n q+(p+2)q-1}(\mathbb{Z}_p, H^*M) &\xrightarrow{i^*} Ext_A^{3,p^n q+(p+2)q-1}(\mathbb{Z}_p, \mathbb{Z}_p) \end{aligned}$$

induced by (1.1). Since $Ext_A^{2,p^n q+(p+2)q-1}(\mathbb{Z}_p, \mathbb{Z}_p) = 0 = Ext_A^{3,p^n q+(p+2)q-1}(\mathbb{Z}_p, \mathbb{Z}_p)$ (cf. [6, Table 8.1]), the above j^* is an isomorphism. Moreover we also know $Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p) = \mathbb{Z}_p \{h_n g_0\}$ (cf. [6, Table 8.1]). Thus we can have that $Ext_A^{3,p^n q+(p+2)q-1}(\mathbb{Z}_p, H^*M) = \mathbb{Z}_p \{j^*(h_n g_0)\}$.

Now observe the following exact sequence

$$\begin{aligned} Ext_A^{2,p^n q+(p+2)q-1}(\mathbb{Z}_p, H^*M) &\xrightarrow{p_*} Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, H^*M) \xrightarrow{i_*} \\ Ext_A^{3,p^n q+(p+2)q}(H^*M, H^*M) &\xrightarrow{j_*} Ext_A^{3,p^n q+(p+2)q-1}(\mathbb{Z}_p, H^*M) \xrightarrow{p_*} \end{aligned}$$

induced by (1.1). Since $Ext_A^{2,p^n q+(p+2)q+r}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $r = -1, 0$ (cf. [2]), we can easily get that $Ext_A^{2,p^n q+(p+2)q-1}(\mathbb{Z}_p, H^*M) = 0$. By virtue of the fact $Ext_A^{3,p^n q+(p+2)q-1}(\mathbb{Z}_p, H^*M) = \mathbb{Z}_p \{j^*(h_n g_0)\}$, we have that the image of the second p_* is $p_* j^*(h_n g_0) = j^* p_*(h_n g_0) = j^* p_*(g_0 h_n) = 0$.

From the facts that

$$Ext_A^{3,p^n q+(p+2)q}(\mathbb{Z}_p, H^*M) \cong \mathbb{Z}_p\{\overline{h_n g_0}\} \cong \mathbb{Z}_p\{j_* \overline{h_n g_0}\}$$

and

$$Ext_A^{3,p^n q+(p+2)q-1}(\mathbb{Z}_p, H^*M) \cong \mathbb{Z}_p\{j^*(h_n g_0)\} \cong \mathbb{Z}_p\{j^* i^* j_* \overline{h_n g_0}\} \cong \mathbb{Z}_p\{j_* j^* i^* \overline{h_n g_0}\},$$

we can easily get that $Ext_A^{3,p^n q+(p+2)q}(H^*M, H^*M) \cong \mathbb{Z}_p\{i_* j_* \overline{h_n g_0}, j^* i^* \overline{h_n g_0}\}$. This shows Proposition 2.3. $\square$

Proposition 2.4 *Let $p \geq 5, n \geq 3$, then we have*

- (1) $i^* d_2(i_* j_* (\overline{h_n g_0})) \neq 0$.
- (2) $d_2(j^* i^* (\overline{h_n g_0})) \neq 0$, where

$$d_2 : Ext_A^{3,p^n q+(p+2)q}(H^*M, H^*M) \rightarrow Ext_A^{5,p^n q+(p+2)q+1}(H^*M, H^*M)$$

is the differential of the Adams spectral sequence.

Proof. (1) From [7, p.488] we know that $d_2(i_*(h_n g_0)) \neq 0$. By Proposition 2.2, $d_2(i_*(h_n g_0)) = d_2(i_* i^* j_* (\overline{h_n g_0})) = d_2(i^* i_* j_* (\overline{h_n g_0})) = i^* d_2(i_* j_* (\overline{h_n g_0}))$. The desired result follows.

(2) Consider the exact sequence

$$Ext_A^{4,p^n q+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p) \xrightarrow{p^*} Ext_A^{5,p^n q+(p+2)q+1}(\mathbb{Z}_p, \mathbb{Z}_p) \xrightarrow{j^*} Ext_A^{5,p^n q+(p+2)q}(\mathbb{Z}_p, H^*M) \xrightarrow{i^*} Ext_A^{5,p^n q+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p)$$

induced by (1.1). We claim that the above j^* is an isomorphism. By virtue of the fact that $Ext_A^{5,p^n q+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ (cf. Proposition 2.1), we see that the above j^* is an epimorphism. Note that $Ext_A^{4,p^n q+(p+2)q}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{g_0 b_{n-1}\}$ (cf. [7, Proposition 2.1]). Since $p^*(g_0 b_{n-1}) = a_0 g_0 b_{n-1} = 0$ (Note: $a_0 g_0 = 0$ by [6, Table 8.2]), so $\ker j^* = \text{imp}^* = 0$, i.e., the above j^* is a monomorphism. The proof of the claim is finished. Since $\alpha_2 b_0 h_n \neq 0 \in Ext_A^{5,p^n q+(p+2)q+1}(\mathbb{Z}_p, \mathbb{Z}_p)$ (cf. [7, Proposition 2.1]), so by the claim we get that $j^*(\alpha_2 b_0 h_n) \neq 0 \in Ext_A^{5,p^n q+(p+2)q}(\mathbb{Z}_p, H^*M)$. Note the fact that $d_2(h_n g_0) = \alpha_2 b_0 h_n \neq 0$, so $j^* d_2(j_* i^* \overline{h_n g_0}) = j_* d_2(j^* i^* \overline{h_n g_0}) \neq 0$ by Proposition 2.2. Thus $d_2(j^* i^* (\overline{h_n g_0})) \neq 0$. $\square$

Proposition 2.5 *Let $p \geq 5, n \geq 3$, then*

$$Ext_A^{3,p^n q+(p+1)q+2}(H^*K, H^*M) = 0.$$

Proof. Consider the exact sequence

$$Ext_A^{3,p^n q+(p+1)q+3}(H^*M, \mathbb{Z}_p) \xrightarrow{j^*} Ext_A^{3,p^n q+(p+1)q+2}(H^*M, H^*M) \xrightarrow{i^*} Ext_A^{3,p^n q+(p+1)q+2}(H^*M, \mathbb{Z}_p).$$

Since the first and third group are zero by the facts $Ext_A^{3,p^n q+(p+1)q+r}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $r = 1, 2, 3$ (cf. [6, Table 8.1]), so the second group is zero.

Look at the exact sequence

$$\begin{aligned} Ext_A^{3,p^n q+pq+2}(\mathbb{Z}_p, \mathbb{Z}_p) &\xrightarrow{i_*} Ext_A^{3,p^n q+pq+2}(H^*M, \mathbb{Z}_p) \xrightarrow{j_*} Ext_A^{3,p^n q+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p) \\ &\xrightarrow{p_*} Ext_A^{4,p^n q+pq+2}(\mathbb{Z}_p, \mathbb{Z}_p) \end{aligned}$$

induced by (1.1). Since we know that $Ext_A^{3,p^n q+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{a_0 h_1 h_n\}$ (cf. [6, Table 8.1]) and $Ext_A^{4,p^n q+pq+2}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{a_0 a_0 h_1 h_n\}$ by Proposition 2.1, so the above p_* is an isomorphism. $im j_* = 0$ since p_* is an isomorphism. $imi_* = 0$ by the fact that $Ext_A^{3,p^n q+pq+2}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ (cf. [6, Table 8.1]). Thus we can have that $Ext_A^{3,p^n q+pq+2}(H^*M, \mathbb{Z}_p) = 0$.

Observe the following exact sequence induced by (1.1)

$$\begin{aligned} Ext_A^{2,p^n q+pq}(\mathbb{Z}_p, \mathbb{Z}_p) &\xrightarrow{p_*} Ext_A^{3,p^n q+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p) \xrightarrow{i_*} Ext_A^{3,p^n q+pq+1}(H^*M, \mathbb{Z}_p) \\ &\xrightarrow{j_*} Ext_A^{3,p^n q+pq}(\mathbb{Z}_p, \mathbb{Z}_p) \xrightarrow{p_*} Ext_A^{4,p^n q+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p). \end{aligned}$$

Since $Ext_A^{2,p^n q+pq}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{h_1 h_n\}$ and $Ext_A^{3,p^n q+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{a_0 h_1 h_n\}$ (cf. [6, Table 8.1]), we know that the first p_* is an isomorphism. Similarly by virtue of the facts that $Ext_A^{3,p^n q+pq}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{b_0 h_n, h_1 b_{n-1}\}$ (cf. [6, Table 8.1]) and $Ext_A^{4,p^n q+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{a_0 b_0 h_n, a_0 h_1 b_{n-1}\}$ (cf. [7, Proposition 2.1]), we get that the second p_* is also an isomorphism. Thus $Ext_A^{3,p^n q+pq+1}(H^*M, \mathbb{Z}_p) = 0$.

Look at the exact sequence

$$\begin{aligned} 0 = Ext_A^{3,p^n q+pq+2}(H^*M, \mathbb{Z}_p) &\xrightarrow{j_*} Ext_A^{3,p^n q+pq+1}(H^*M, H^*M) \xrightarrow{i_*} \\ &Ext_A^{3,p^n q+pq+1}(H^*M, \mathbb{Z}_p) = 0 \end{aligned}$$

induced by (1.1). It is easy to get that the second group is zero.

At last consider the following exact sequence

$$\begin{aligned} 0 = Ext_A^{3,p^n q+(p+1)q+2}(H^*M, H^*M) &\xrightarrow{i'_*} Ext_A^{3,p^n q+(p+1)q+2}(H^*K, H^*M) \\ &\xrightarrow{j'_*} Ext_A^{3,p^n q+pq+1}(H^*M, H^*M) = 0 \end{aligned}$$

induced by (1.2). The desired result follows. $\square$

Proposition 2.6 *Let $p \geq 5, n \geq 3$, then*

$$Ext_A^{2,p^n q+(p+1)q+1}(H^*K, \mathbb{Z}_p) \cong \mathbb{Z}_p\{\beta_* i'_* i_*(h_n)\},$$

where $\beta_* : Ext_A^{1,p^n q}(H^*K, \mathbb{Z}_p) \longrightarrow Ext_A^{2,p^n q+(p+1)q+1}(H^*K, \mathbb{Z}_p)$ is the connecting homomorphism induced by $\beta : \Sigma^{(p+1)q}K \longrightarrow K$.

Proof. Look at the exact sequence

$$\begin{aligned} Ext_A^{1,p^n q+pq-1}(\mathbb{Z}_p, \mathbb{Z}_p) &\xrightarrow{p_*} Ext_A^{2,p^n q+pq}(\mathbb{Z}_p, \mathbb{Z}_p) \xrightarrow{i_*} Ext_A^{2,p^n q+pq}(H^*M, \mathbb{Z}_p) \\ &\xrightarrow{j_*} Ext_A^{2,p^n q+pq-1}(\mathbb{Z}_p, \mathbb{Z}_p) \end{aligned}$$

induced by (1.1). Since the first group and the fourth group are zero, so the above i_* is an isomorphism. Thus we can see that $Ext_A^{2,p^n q+pq}(H^*M, \mathbb{Z}_p) \cong \mathbb{Z}_p\{i_*(h_1 h_n)\}$ by the fact that $Ext_A^{2,p^n q+pq}(\mathbb{Z}_p, \mathbb{Z}_p) \cong \mathbb{Z}_p\{h_1 h_n\}$.

At last observe the following exact sequence

$$\begin{array}{ccc} Ext_A^{2,p^ng+(p+1)q+1}(H^*M, \mathbb{Z}_p) & \xrightarrow{i'_*} & Ext_A^{2,p^ng+(p+1)q+1}(H^*K, \mathbb{Z}_p) \xrightarrow{j'_*} \\ & & Ext_A^{2,p^ng+pq}(H^*M, \mathbb{Z}_p) \xrightarrow{\alpha_*} Ext_A^{3,p^ng+(p+1)q+1}(H^*M, \mathbb{Z}_p) \end{array}$$

induced by (1.2). Since the first group is zero by $Ext_A^{2,p^ng+(p+1)q+r}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $r = 0, 1$ (cf. [2]) and the fourth group is zero by $Ext_A^{3,p^ng+(p+1)q+t}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $t = 0, 1$ (cf. [6, Table 8.1]), then the above j'_* is an isomorphism. Thus we can have that $Ext_A^{2,p^ng+(p+1)q+1}(H^*K, \mathbb{Z}_p)$ has a unique generator Δ , which satisfies $j'_*(\Delta) = i_*(h_1h_n)$. From [5, (5.4)], we have that $j'\beta i'i \in [\sum^{pq-1} S, M]$ is represented by $i_*(h_1) \in Ext_A^{1,pq}(H^*M, \mathbb{Z}_p)$ in the Adams spectral sequence. It follows that $(j'\beta i'i)_*(h_n) = i_*(h_1h_n) = j'_*(\Delta)$. Note the fact that j'_* is an isomorphism. It is easy to get that $\beta_*i'_*i_*(h_n) = \Delta$. Therefore this completes the proof of the proposition. $\square$

Proposition 2.7 *Let $p \geq 5, n \geq 3$, then*

$$Ext_A^{2,p^ng+(p+1)q+1}(H^*K, H^*M) \cong \mathbb{Z}_p\{\beta_*i'_*(\tilde{h}_n)\}$$

where $\tilde{h}_n \in Ext_A^{1,p^ng}(H^*M, H^*M)$ is the unique generator of $Ext_A^{1,p^ng}(H^*M, H^*M)$ and satisfies $i^*(\tilde{h}_n) = i_*(h_n)$.

Proof. Consider the exact sequence

$$\begin{array}{ccccc} Ext_A^{2,p^ng+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p) & \xrightarrow{i_*} & Ext_A^{2,p^ng+pq+1}(H^*M, \mathbb{Z}_p) & \xrightarrow{j_*} & Ext_A^{2,p^ng+pq}(\mathbb{Z}_p, \mathbb{Z}_p) \\ & & & & \xrightarrow{p_*} Ext_A^{3,p^ng+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p). \end{array}$$

Since $Ext_A^{2,p^ng+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$, $imi_* = 0$. Since $Ext_A^{2,p^ng+pq}(\mathbb{Z}_p, \mathbb{Z}_p) = \mathbb{Z}_p\{h_1h_n\}$ and $Ext_A^{3,p^ng+pq+1}(\mathbb{Z}_p, \mathbb{Z}_p) = \mathbb{Z}_p\{a_0h_1h_n\}$ (cf. [6, Table 8.1]), then the above p_* is an isomorphism, and then $imj_* = 0$. Thus $Ext_A^{2,p^ng+pq+1}(H^*M, \mathbb{Z}_p) = 0$.

Look at the exact sequence

$$\begin{array}{ccc} Ext_A^{2,p^ng+(p+1)q+2}(H^*M, \mathbb{Z}_p) & \xrightarrow{i'_*} & Ext_A^{2,p^ng+(p+1)q+2}(H^*K, \mathbb{Z}_p) \xrightarrow{j'_*} \\ & & Ext_A^{2,p^ng+pq+1}(H^*M, \mathbb{Z}_p) = 0 \end{array}$$

induced by (1.2). Since we know that $Ext_A^{2,p^ng+(p+1)q+2}(H^*M, \mathbb{Z}_p) = 0$ by the facts that $Ext_A^{2,p^ng+(p+1)q+r}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $r = 1, 2$ and $Ext_A^{2,p^ng+pq+1}(H^*M, \mathbb{Z}_p) = 0$, then $Ext_A^{2,p^ng+(p+1)q+2}(H^*K, \mathbb{Z}_p) = 0$.

Observe the following exact sequence

$$Ext_A^{3,p^ng+(p+1)q+2}(H^*M, \mathbb{Z}_p) \xrightarrow{i'_*} Ext_A^{3,p^ng+(p+1)q+2}(H^*K, \mathbb{Z}_p) \xrightarrow{j'_*} Ext_A^{3,p^ng+pq+1}(H^*M, \mathbb{Z}_p)$$

induced by (1.2). From the proof of Proposition 2.5 we know that the first group and the third group are zero. So the middle group is zero.

At last, look at the following exact sequence

$$\begin{aligned} Ext_A^{2,p^ng+(p+1)q+2}(H^*K, \mathbb{Z}_p) &\xrightarrow{j^*} Ext_A^{2,p^ng+(p+1)q+1}(H^*K, H^*M) \xrightarrow{i^*} \\ &Ext_A^{2,p^ng+(p+1)q+1}(H^*K, \mathbb{Z}_p) \xrightarrow{p^*} Ext_A^{3,p^ng+(p+1)q+2}(H^*K, \mathbb{Z}_p) \end{aligned}$$

induced by (1.1). Since the first group and the fourth group are zero, then the above i^* is an isomorphism. Notice that $Ext_A^{2,p^ng+(p+1)q+1}(H^*K, \mathbb{Z}_p) \cong \mathbb{Z}_p\{\beta_*i'_*i_*(h_n)\}$. Thus we can easily have that there exists an element $\bar{\Delta} \in Ext_A^{2,p^ng+(p+1)q+1}(H^*K, H^*M)$ such that $Ext_A^{2,p^ng+(p+1)q+1}(H^*K, H^*M) \cong \mathbb{Z}_p\{\bar{\Delta}\}$ and $i^*(\bar{\Delta}) = \beta_*i'_*i_*(h_n)$. Since $i^*(\tilde{h}_n) = i_*(h_n)$, $i^*(\bar{\Delta}) = \beta_*i'_*i_*(h_n) = \beta_*i'_*i^*(\tilde{h}_n) = i^*\beta_*i'_*(\tilde{h}_n)$. Thus we have that $\bar{\Delta} = \beta_*i'_*(\tilde{h}_n)$ by the fact that i^* is an isomorphism. Thus this completes the proof of the proposition. $\square$

3 Proofs of the main theorems

Let

$$\begin{array}{ccccc} \dots & \xrightarrow{\bar{a}_2} & \Sigma^{-2}E_2 & \xrightarrow{\bar{a}_1} & \Sigma^{-1}E_1 & \xrightarrow{\bar{a}_0} & S \\ & & \downarrow \bar{b}_2 & & \downarrow \bar{b}_1 & & \downarrow \bar{b}_0 \\ & & \Sigma^{-2}KG_2 & & \Sigma^{-1}KG_1 & & KG_0 = K\mathbb{Z}_p \end{array} \quad (3.1)$$

be the minimal Adams resolution of S satisfying the following.

(1) $E_s \xrightarrow{\bar{b}_s} KG_s \xrightarrow{\bar{c}_s} E_{s+1} \xrightarrow{\bar{a}_s} \Sigma E_s$ are cofibrations for all $s \geq 0$ which induce short exact sequences in $\mathbb{Z}_p$ -cohomology.

(2) KG_s is a wedge sum of suspensions of Eilenberg-MacLane spectra of type $K\mathbb{Z}_p$.

(3) $\pi_t KG_s$ are the $E_1^{s,t}$ -terms, $(\bar{b}_s \bar{c}_{s-1})_* : \pi_t KG_{s-1} \rightarrow \pi_t KG_s$ are the $d_1^{s-1,t}$ -differentials of the Adams spectral sequence and $\pi_t KG_s \cong Ext_A^{s,t}(\mathbb{Z}_p, \mathbb{Z}_p)$ (cf. [9, p.180]). Then

$$\begin{array}{ccccc} \dots & \xrightarrow{\bar{a}_2 \wedge 1_W} & \Sigma^{-2}E_2 \wedge W & \xrightarrow{\bar{a}_1 \wedge 1_W} & \Sigma^{-1}E_1 \wedge W & \xrightarrow{\bar{a}_0 \wedge 1_W} & W \\ & & \downarrow \bar{b}_2 \wedge 1_W & & \downarrow \bar{b}_1 \wedge 1_W & & \downarrow \bar{b}_0 \wedge 1_W \\ & & \Sigma^{-2}KG_2 \wedge W & & \Sigma^{-1}KG_1 \wedge W & & KG_0 \wedge W \end{array} \quad (3.2)$$

is an Adams resolution of arbitrary finite spectrum W .

From [10, p.204-206], the Moore spectrum M is a commutative ring spectrum with multiplication $m_M : M \wedge M \rightarrow M$ and there is $\bar{m}_M : \Sigma M \rightarrow M \wedge M$ such that

$$\begin{aligned} m_M(i \wedge 1_M) &= 1_M, & (j \wedge 1_M)\bar{m}_M &= 1_M, \\ m_M\bar{m}_M &= 0, & (i \wedge 1_M)m_M + \bar{m}_M(j \wedge 1_M) &= 1_{M \wedge M}, \\ m_M T &= -m_M, & T\bar{m}_M &= \bar{m}_M, \\ m_M(1_M \wedge i) &= -1_M, & (1_M \wedge j)\bar{m}_M &= 1_M, \end{aligned}$$

where $T : M \wedge M \rightarrow M \wedge M$ is the switching map.

A spectrum X is called an M -module spectrum if $p \wedge 1_X = 0$, and consequently, the cofibration $X \xrightarrow{p \wedge 1_X} X \xrightarrow{i \wedge 1_X} M \wedge X \xrightarrow{j \wedge 1_X} \Sigma X$ split, i.e., there is a homotopy

equivalence $M \wedge X = X \vee \Sigma X$ and there are maps $m_X : M \wedge X \longrightarrow X$, $\bar{m}_X : \Sigma X \longrightarrow M \wedge X$ satisfying $m_X(i \wedge 1_X) = 1_X$, $(j \wedge 1_X)\bar{m}_X = 1_X$, $m_X\bar{m}_X = 0$ and $\bar{m}_X(j \wedge 1_X) + (i \wedge 1_X)m_X = 1_{M \wedge X}$. The M -module actions m_X , $\bar{m}_X$ are called associative if $m_X(1_M \wedge m_X) = -m_X(m_X \wedge 1_X)$ and $(1_M \wedge \bar{m}_M)\bar{m}_X = (\bar{m}_M \wedge 1_X)\bar{m}_X$.

Let X and X' be M -module spectra. Then we define a homomorphism $d : [\Sigma^s X', X] \rightarrow [\Sigma^{s+1} X', X]$ by $d(f) = m_X(1_M \wedge f)\bar{m}_{X'}$ for $f \in [\Sigma^s X', X]$. This operation d is called a derivation (of maps between M -module spectra) which has the following properties:

Lemma 3.1^[10, Theorem 2.2] (1) d is a derivative: $d(fg) = fd(g) + (-1)^{|g|}d(f)g$ for $f \in [\Sigma^s X', X]$, $g \in [\Sigma^t X'', X']$, where X , X' , X'' are M -module spectra.

(2) Let W' , W be arbitrary spectra and $h \in [\Sigma^r W', W]$. Then

$$d(h \wedge f) = (-1)^{|h|}h \wedge d(f)$$

for $f \in [\Sigma^s X', X]$.

(3) $d^2 = 0 : [\Sigma^s X', X] \longrightarrow [\Sigma^{s+2} X', X]$ for associative spectra X' , X . $\square$

From [10, (3.4)], K is an M -module spectrum, i.e., there are M -module actions $m_K : K \wedge M \longrightarrow K$, $\bar{m}_K : \Sigma K \longrightarrow K \wedge M$ satisfying

$$m_K(1_K \wedge i) = 1_K, (1_K \wedge j)\bar{m}_K = 1_K, m_K\bar{m}_K = 0, (1_K \wedge i)m_K + (1_K \wedge j)\bar{m}_K = 1_{K \wedge M}.$$

Moreover, from [10, (2.6)] and [10, (3.7)] we have that $d(ij) = -1_M$, $d(\alpha) = 0$, $d(i') = 0$, $d(j') = 0$ and $d(\beta) = 0$.

Remark 3.1 In this paper, all the notations are the same as those of [7].

Let L be the cofiber of $\alpha_1 = j\alpha_i : \Sigma^{q-1}S \rightarrow S$ and K' be the cofiber of $\alpha i : \Sigma^q S \rightarrow M$ given by the following two cofibrations:

$$\Sigma^{q-1}S \xrightarrow{\alpha_1} S \xrightarrow{i''} L \xrightarrow{j''} \Sigma^q S (\text{see}[7, (2.3)]), \quad (3.3)$$

$$\Sigma^q S \xrightarrow{\alpha i} M \xrightarrow{v} K' \xrightarrow{y} \Sigma^{q+1} S (\text{see}[7, (2.4)]). \quad (3.4)$$

Let $\alpha' = \alpha_1 \wedge 1_K$. Consider the following two commutative diagram of 3×3 in the stable homotopy category:

$$\begin{array}{ccccccc} \Sigma M & & \xrightarrow{v} & & \Sigma K' & & \xrightarrow{1_{K'} \wedge p} & & \Sigma K' \\ & \searrow (v \wedge 1_M)\bar{m}_M & & \nearrow 1_{K'} \wedge j & & \searrow y & & \nearrow z & \\ & & K' \wedge M & & & & \Sigma^{q+2} S & & \\ & \nearrow 1_{K'} \wedge i & & \searrow \pi & & \nearrow jj' & & \searrow \alpha i & \\ K' & & \xrightarrow{x} & & K & & \xrightarrow{j' \alpha'} & & \Sigma^2 M \end{array}$$

and

$$\begin{array}{ccccccc} M & & \xrightarrow{(i'' \wedge 1_K)i'} & & L \wedge K & & \xrightarrow{j'' \wedge 1_K} & & \Sigma^q K \\ & \searrow i' & & \nearrow i'' \wedge 1_K & & \searrow \bar{r} & & \nearrow \pi & \\ & & K & & & & \Sigma^q K' \wedge M & & \\ & \nearrow \alpha' & & \searrow j' & & \nearrow (v \wedge 1_M)\bar{m}_M & & \searrow \varepsilon & \\ \Sigma^{q-1} K & & \xrightarrow{j' \alpha'} & & \Sigma^{q+1} M & & \xrightarrow{\alpha} & & \Sigma M \end{array}$$

By the above two commutative diagram of 3×3 in the stable homotopy category, we easily have the following two lemmas.

Lemma 3.2 *There exist three cofibrations*

$$K' \xrightarrow{x} K \xrightarrow{jj'} \Sigma^{q+2} S \xrightarrow{z} \Sigma K', \quad (3.5)$$

$$\Sigma^{-1} K \xrightarrow{j'\alpha'} \Sigma M \xrightarrow{(v \wedge 1_M) \bar{m}_M} K' \wedge M \xrightarrow{\pi} K, \quad (3.6)$$

$$M \xrightarrow{(i'' \wedge 1_K) i'} L \wedge K \xrightarrow{\bar{r}} \Sigma^q K' \wedge M \xrightarrow{\varepsilon} \Sigma M. \quad (3.7)$$

□

Lemma 3.3 $\varepsilon(v \wedge 1_M) \bar{m}_M = \alpha$, $\bar{r}(i'' \wedge 1_K) = (v \wedge 1_M) \bar{m}_M j'$, $\pi \bar{r} = j'' \wedge 1_K$, $\varepsilon(1_{K'} \wedge i) v j' = -2j' \alpha'$. □

From [11, p.434], there are $\bar{\Delta} \in [\Sigma^{-1-1} L \wedge K, K]$ and $\tilde{\Delta} \in [\Sigma^{-1} K, L \wedge K]$ satisfying $\bar{\Delta}(i'' \wedge 1_K) = (j'' \wedge 1_K) \tilde{\Delta} = i' j' \in [\Sigma^{-q-1} K, K]$ and $j j' \bar{\Delta} = 0$. From [6, p.484], there is $\bar{\Delta}_{K'} \in [\Sigma^{-q-1} L \wedge K, K']$ such that $\bar{\Delta}_{K'}(i'' \wedge 1_K) = v j' \in [\Sigma^{-q-1} K, K']$ and $\bar{\Delta}(i'' \wedge 1_K) = (j'' \wedge 1_K) \tilde{\Delta} = i' j'$.

Lemma 3.4 $\bar{\Delta}_{K'} = (1_{K'} \wedge j) \bar{r}$.

Proof. From Lemma 3.3 we have

$$(1_{K'} \wedge j) \bar{r}(i'' \wedge 1_K) = (1_{K'} \wedge j)(v \wedge 1_M) \bar{m}_M j' = (v \wedge 1_{S^0})(1_M \wedge j) \bar{m}_M j' = v j' = \bar{\Delta}_{K'}(i'' \wedge 1_K),$$

which shows that $(1_{K'} \wedge j) \bar{r} = \bar{\Delta}_{K'} + \bar{g}(j'' \wedge 1_K)$ for some $\bar{g} \in [K, \Sigma K']$

Consider the exact sequence induced by (3.4)

$$[K, \Sigma^{q+1} S] \xrightarrow{(\alpha i)^*} [K, \Sigma M] \xrightarrow{v_*} [K, \Sigma K'] \xrightarrow{y_*} [K, \Sigma^{q+2} S] \xrightarrow{(\alpha i)^*} [K, \Sigma^2 M].$$

From the proof of [7, Proposition 2.18], we know that $[K, \Sigma M] = 0$. So $im v_* = 0$. Since $[K, \Sigma^{q+2} S] \cong \mathbb{Z}_p \{jj'\}$, $(\alpha i)_*(jj') = \alpha i j j' \neq 0$. Thus we have $im y_* = 0$ and $[K, \Sigma K'] = 0$. Then we have $(1_{K'} \wedge j) \bar{r} = \bar{\Delta}_{K'}$. □

Lemma 3.5 [7, lemma 3.3 and (3.4)] *Let $p \geq 5, n \geq 3$, then there exists an element $\eta'_{n,2} \in [\Sigma^{p^n q+q} K, E_2 \wedge K]$ such that*

$$(\bar{b}_2 \wedge 1_K) \eta'_{n,2} = h_0 h_n \wedge 1_K \in [\Sigma^{p^n q+q} K, K G_2 \wedge K], (1_{E_2} \wedge \alpha') \eta'_{n,2} = 0,$$

where $h_0 h_n \in \pi_{p^n q+q} K G_2 \cong Ext_A^{2,p^n q+q}(\mathbb{Z}_p, \mathbb{Z}_p)$ and $\alpha' = j \alpha i \wedge 1_K \in [\Sigma^{q-1} K, K]$. There also exists an element $f_2 \in [\Sigma^{p^n q+(p+2)q+3} M, E_5 \wedge L \wedge K]$ such that

$$(1_{E_2} \wedge (i'' \wedge 1_K) \beta) \eta'_{n,2} i' = (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{L \wedge K}) f_2.$$

□

Corollary 3.1 *For $f_2 \in [\Sigma^{p^n q+(p+2)q+3} M, E_5 \wedge L \wedge K]$ which is given in Lemma 3.5, we have*

$$(1_{E_4} \wedge \varepsilon(1_{K'} \wedge i j)) (\bar{a}_4 \wedge 1_{K' \wedge M}) (1_{E_5} \wedge \bar{r}) d(f_2 i j) = 0. \quad (3.8)$$

Proof. From [7], we have [7, (3.6)] that $(\bar{a}_4 \wedge 1_M)(1_{E_5} \wedge \varepsilon(1_{K'} \wedge i)\bar{\Delta}_{K'})d(f_2ij) = 0$. Here f_2 is given in [7, (3.4)].

By [10, (1.7)], we have that $(\bar{a}_4 \wedge 1_M)(1_{E_5} \wedge \varepsilon(1_{K'} \wedge i))(1_{E_5} \wedge \bar{\Delta}_{K'})d(f_2ij) = 0$. By lemma 3.4, we have

$$(\bar{a}_4 \wedge 1_M)(1_{E_5} \wedge \varepsilon(1_{K'} \wedge i))(1_{E_5} \wedge (1_{K'} \wedge j)\bar{r})d(f_2ij) = 0.$$

By [10, (1.7)], it follows that

$$(\bar{a}_4 \wedge 1_M)(1_{E_5} \wedge \varepsilon(1_{K'} \wedge i))(1_{E_5} \wedge (1_{K'} \wedge j))(1_{E_5} \wedge \bar{r})d(f_2ij) = 0.$$

Thus

$$(\bar{a}_4 \wedge 1_M)(1_{E_5} \wedge \varepsilon(1_{K'} \wedge ij))(1_{E_5} \wedge \bar{r})d(f_2ij) = 0.$$

By [10, (1.7)], the corollary follows. $\square$

Let W be the cofibre of $\varepsilon(1_{K'} \wedge ij) : \Sigma^{q-2}K' \wedge M \longrightarrow M$ given by the cofibration

$$\Sigma^{q-2}K' \wedge M \xrightarrow{\varepsilon(1_{K'} \wedge ij)} M \xrightarrow{w_4} W \xrightarrow{u_4} \Sigma^{q-1}K' \wedge M. \quad (3.9)$$

Lemma 3.6 *There exists $f' \in [\Sigma^{p^n q + (p+2)q+1}M, E_4 \wedge W]$ such that $(\bar{a}_2\bar{a}_3 \wedge 1_K)(1_{E_4} \wedge \pi u_4)f' = 0$.*

Proof. By (3.8) and (3.9), we have that

$$(\bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r})d(f_2ij) = (1_{E_4} \wedge u_4)f' \quad (3.10)$$

with $f' \in [\Sigma^{p^n q + (p+2)q+1}M, E_4 \wedge W]$ and by composing $(\bar{a}_2\bar{a}_3 \wedge 1_K)(1_{E_4} \wedge \pi)$ on (3.10) we have

$$(\bar{a}_2\bar{a}_3 \wedge 1_K)(1_{E_4} \wedge \pi u_4)f' = (\bar{a}_2\bar{a}_3\bar{a}_4 \wedge 1_K)(1_{E_5} \wedge \pi\bar{r})d(f_2ij). \quad (3.11)$$

By composing ij on [7, (3.4)], we have

$$(1_{E_2} \wedge (i'' \wedge 1_K)\beta)\eta'_{n,2}i'ij = (\bar{a}_2\bar{a}_3\bar{a}_4 \wedge 1_{L \wedge K})f_2ij \quad (3.12)$$

with $\eta'_{n,2} \in [\Sigma^{p^n q + q}K, E_2 \wedge K]$.

Notice that $d(1_K) = 0$ and $d(\beta) = 0$. Then by applying the derivation d on (3.12) we have

$$(1_{E_2} \wedge (i'' \wedge 1_K)\beta)d(\eta'_{n,2}i'ij) = (\bar{a}_2\bar{a}_3\bar{a}_4 \wedge 1_{L \wedge K})d(f_2ij). \quad (3.13)$$

Notice that $\pi\bar{r} = j'' \wedge 1_K$. By composing $(1_{E_2} \wedge \pi\bar{r})$ on (3.13) we have

$$(\bar{a}_2\bar{a}_3\bar{a}_4 \wedge 1_K)(1_{E_5} \wedge \pi\bar{r})d(f_2ij) = 0 \quad (3.14)$$

and by (3.11), (3.14) we get

$$(\bar{a}_2\bar{a}_3 \wedge 1_K)(1_{E_4} \wedge \pi u_4)f' = 0. \quad (3.15)$$

Thus the Lemma is proved. $\square$

Let U be the cofibre of $\pi u_4 : W \longrightarrow \Sigma^{q-1}K$ given by the cofibration

$$W \xrightarrow{\pi u_4} \Sigma^{q-1}K \xrightarrow{w_5} U \xrightarrow{u_5} \Sigma W. \quad (3.16)$$

Lemma 3.7 *w_5 induces zero homomorphism in $\mathbb{Z}_p$ -cohomology.*

Proof. Consider the following homomorphism induced by w_5 :

$$w_5^* : H^*U \longrightarrow H^{*+q-1}K.$$

From the cellular structures of U and K , we can have that

$$H^t K = \begin{cases} \mathbb{Z}_p, & t = 0, 1, q+1, q+2; \\ 0, & \text{others,} \end{cases}$$

and the top cell of U has degree $2q+1$. It easily follows that w_5^* must be a zero homomorphism in $\mathbb{Z}_p$ -cohomology. $\square$

Lemma 3.8 *There exist three homotopy elements $f'_2 \in [\Sigma^{p^n q + (p+2)q} M, E_2 \wedge U]$, $f'_3 \in [\Sigma^{p^n q + (p+2)q+1} M, E_3 \wedge U]$ and $g_2 \in [\Sigma^{p^n q + (p+2)q} M, KG_2 \wedge W]$ such that*

$$(\bar{a}_2 \bar{a}_3 \wedge 1_W) f' = (1_{E_2} \wedge u_5) f'_2, f'_2 = (\bar{a}_2 \wedge 1_U) f'_3$$

and

$$(1_{E_3} \wedge u_4)(\bar{a}_3 \wedge 1_W) f' = -(1_{E_3} \wedge u_4 u_5) f'_3 + (1_{E_3} \wedge u_4)(\bar{c}_2 \wedge 1_W) g_2.$$

Proof. From (3.15) and (3.16), we have that

$$(\bar{a}_2 \bar{a}_3 \wedge 1_W) f' = (1_{E_2} \wedge u_5) f'_2 \quad (3.17)$$

with $f'_2 \in [\Sigma^{p^n q + (p+2)q} M, E_2 \wedge U]$.

By (3.17) and (3.2) we have $(\bar{b}_2 \wedge 1_W)(1_{E_2} \wedge u_5) f'_2 = (\bar{b}_2 \wedge 1_W)(\bar{a}_2 \bar{a}_3 \wedge 1_W) f' = 0$. Thus

$$(1_{KG_2} \wedge u_5)(\bar{b}_2 \wedge 1_U) f'_2 = 0. \quad (3.18)$$

By (3.18), (3.16) and the fact that w_5 induces zero homomorphism in $\mathbb{Z}_p$ -cohomology (see lemma 3.7), we have

$$(\bar{b}_2 \wedge 1_U) f'_2 = (1_{KG_2} \wedge w_5) g = 0 \quad (3.19)$$

with $g \in [\Sigma^{p^n q + (p+1)q+1} M, KG_2 \wedge K]$, so by (3.2) we obtain

$$f'_2 = (\bar{a}_2 \wedge 1_U) f'_3 \quad (3.20)$$

with $f'_3 \in [\Sigma^{p^n q + (p+2)q+1} M, E_3 \wedge U]$. By [10, (1.7)], from (3.20) and (3.17) we have $(\bar{a}_2 \bar{a}_3 \wedge 1_W) f' = -(\bar{a}_2 \wedge 1_W)(1_{E_3} \wedge u_5) f'_3$. Then we have

$$(\bar{a}_3 \wedge 1_W) f' = -(1_{E_3} \wedge u_5) f'_3 + (\bar{c}_2 \wedge 1_W) g_2 \quad (3.21)$$

with $g_2 \in [\Sigma^{p^n q + (p+2)q} M, KG_2 \wedge W]$. By composing $(1_{E_3} \wedge u_4)$ on (3.21), we have

$$(1_{E_3} \wedge u_4)(\bar{a}_3 \wedge 1_W)f' = -(1_{E_3} \wedge u_4 u_5)f'_3 + (1_{E_3} \wedge u_4)(\bar{c}_2 \wedge 1_W)g_2. \quad (3.22)$$

We finishes the proof of the lemma. $\square$

Lemma 3.9 *the cofibre of $\varepsilon(1_{K'} \wedge i)v : \Sigma^q M \longrightarrow \Sigma M$ is U given by the cofibration*

$$\Sigma^q M \xrightarrow{\varepsilon(1_{K'} \wedge i)v} \Sigma M \xrightarrow{w_6} U \xrightarrow{u_6} \Sigma^{q+1} M. \quad (3.23)$$

There exist two relations that

$$u_4 u_5 = (v \wedge 1_M) \bar{m}_M u_6$$

and

$$\varepsilon(1_{K'} \wedge i j)(v \wedge 1_M) \bar{m}_M = \varepsilon(1_{K'} \wedge i)v.$$

Proof. By the three cofibrations (3.6), (3.9), and (3.16), we can get the following commutative diagram (3.24) of 3×3 lemma in stable homotopy category(cf. [12, p. 292-293]).

$$\begin{array}{ccccccc} & W & \xrightarrow{\pi u_4} & & \Sigma^{q-1} K & \xrightarrow{j' \alpha'} & \Sigma^{q+1} M \\ u_4 \searrow & & & \nearrow \pi & \searrow w_5 & & \nearrow u_6 \\ v \wedge 1_M \bar{m}_M \nearrow & \Sigma^{q-1} K' \wedge M & & \searrow \varepsilon(1_{K'} \wedge i j) & \nearrow w_6 & U & \searrow u_5 \\ & \xrightarrow{\varepsilon(1_{K'} \wedge i)v} & & \Sigma M & \xrightarrow{w_4} & \Sigma W & \nearrow u_4 \\ & & & & & & \Sigma^q K' \wedge M. \end{array}$$

By the commutative diagram (3.24), Lemma 3.8 follows. $\square$

Lemma 3.10 *With notation as above. We have*

$$\begin{aligned} & (\bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r})d(f_2 i j) \\ = & (1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M u_6) f'_3 - (\bar{c}_2 \wedge 1_{K' \wedge M})(1_{KG_2} \wedge u_4) g_2. \end{aligned}$$

Proof. By (3.22), [10, (1.7)] and the relation $u_4 u_5 = (v \wedge 1_M) \bar{m}_M u_6$ (see Lemma 3.9), we have

$$(\bar{a}_3 \wedge 1_{K' \wedge M})(1_{E_4} \wedge u_4) f' = (1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M u_6) f'_3 - (\bar{c}_2 \wedge 1_{K' \wedge M})(1_{KG_2} \wedge u_4) g_2. \quad (3.25)$$

By composing $(\bar{a}_3 \wedge 1_{K' \wedge M})$ on (3.10), we have

$$(\bar{a}_3 \wedge 1_{K' \wedge M})(1_{E_4} \wedge u_4) f' = (\bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r})d(f_2 i j). \quad (3.26)$$

Combining (3.25) and (3.26) yields

$$\begin{aligned} & (\bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r})d(f_2 i j) \\ = & (1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M u_6) f'_3 - (\bar{c}_2 \wedge 1_{K' \wedge M})(1_{KG_2} \wedge u_4) g_2. \end{aligned} \quad (3.27)$$

Thus we complete the proof of this lemma. $\square$

Lemma 3.11 *There exist two elements $f'_4 \in [\Sigma^{p^n q + (p+2)q+1} M, E_3 \wedge K]$ and $f'_5 \in [\Sigma^{p^n q + (p+2)q+1} M, E_3 \wedge K' \wedge M]$ such that*

$$(1_{E_3} \wedge u_6) f'_3 = (1_{E_3} \wedge j') f'_4$$

and

$$f'_4 = (1_{E_3} \wedge \pi) f'_5.$$

Proof. By Lemma 3.3, we have $\alpha = \varepsilon(v \wedge 1_M) \bar{m}_M$. Then

$$\begin{aligned} & (1_{E_3} \wedge \alpha u_6) f'_3 \\ = & (1_{E_3} \wedge \varepsilon(v \wedge 1_M) \bar{m}_M u_6) f'_3, \quad \text{since } u_4 u_5 = (v \wedge 1_M) \bar{m}_M u_6 \\ = & (1_{E_3} \wedge \varepsilon u_4 u_5) f'_3 \\ = & (1_{E_3} \wedge \varepsilon)(1_{E_3} \wedge u_4 u_5) f'_3, \quad \text{by (3.22)} \\ = & (1_{E_3} \wedge \varepsilon)[(1_{E_3} \wedge u_4)(\bar{c}_2 \wedge 1_W) g_2 - (1_{E_3} \wedge u_4)(\bar{a}_3 \wedge 1_W) f'] \\ = & (1_{E_3} \wedge \varepsilon u_4)(\bar{c}_2 \wedge 1_W) g_2 - (1_{E_3} \wedge \varepsilon u_4)(\bar{a}_3 \wedge 1_W) f' \\ = & (\bar{c}_2 \wedge 1_M)(1_{KG_2} \wedge \varepsilon u_4) g_2 - (1_{E_3} \wedge \varepsilon u_4)(\bar{a}_3 \wedge 1_W) f', \quad \text{since } 1_{KG_2} \wedge \varepsilon \simeq 0 \\ = & -(1_{E_3} \wedge \varepsilon u_4)(\bar{a}_3 \wedge 1_W) f' \\ = & (1_{E_3} \wedge \varepsilon)(\bar{a}_3 \wedge 1_{K' \wedge M})(1_{E_4} \wedge u_4) f', \quad \text{by (3.26)} \\ = & (1_{E_3} \wedge \varepsilon)(\bar{a}_3 \wedge 1_{K' \wedge M})(\bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r}) d(f_2 i j) \\ = & (\bar{a}_3 \bar{a}_4 \wedge 1_M)(1_{E_5} \wedge \varepsilon \bar{r}) d(f_2 i j), \quad \text{by (3.7)} \\ = & 0. \end{aligned}$$

Hence, by (1.2) we have

$$(1_{E_3} \wedge u_6) f'_3 = (1_{E_3} \wedge j') f'_4 \tag{3.28}$$

with $f'_4 \in [\Sigma^{p^n q + (p+2)q+1} M, E_3 \wedge K]$.

Similarly, by Lemma 3.3 we have $\varepsilon(1_{K'} \wedge i) v j' = -2j' \alpha'$. Then we have

$$\begin{aligned} & -2(1_{E_3} \wedge j' \alpha') f'_4 \\ = & (1_{E_3} \wedge \varepsilon(1_{K'} \wedge i) v j') f'_4 \\ = & (1_{E_3} \wedge \varepsilon(1_{K'} \wedge i) v)(1_{E_3} \wedge j') f'_4, \quad \text{by (3.28)} \\ = & (1_{E_3} \wedge \varepsilon(1_{K'} \wedge i) v)(1_{E_3} \wedge u_6) f'_3, \quad \text{by (3.24)} \\ = & (1_{E_3} \wedge \varepsilon(1_{K'} \wedge i j))(v \wedge 1_M) \bar{m}_M (1_{E_3} \wedge u_6) f'_3 \\ = & (1_{E_3} \wedge \varepsilon(1_{K'} \wedge i j))(1_{E_3} \wedge v \wedge 1_M \bar{m}_M u_6) f'_3, \quad \text{by (3.24)} \\ = & (1_{E_3} \wedge \varepsilon(1_{K'} \wedge i j))(1_{E_3} \wedge u_4 u_5) f'_3, \quad \text{by (3.9)} \\ = & (1_{E_3} \wedge \varepsilon(1_{K'} \wedge i j) u_4 u_5) f'_3 \\ = & 0. \end{aligned}$$

Thus, by (3.6) we have

$$f'_4 = (1_{E_3} \wedge \pi) f'_5 \tag{3.29}$$

with $f'_5 \in [\Sigma^{p^n q + (p+2)q+1} M, E_3 \wedge K' \wedge M]$. This completes the proof of Lemma 3.1 $\square$

Lemma 3.12 *For the above $f'_5 \in [\Sigma^{p^n q + (p+2)q+1} M, E_3 \wedge K' \wedge M]$, we have*

$$(\bar{b}_3 \wedge 1_{K' \wedge M}) f'_5 = 0.$$

Proof. The proof will be given later. □

Now we give the proof of Theorem I.

Proof of Theorem I. From Lemma 3.12, we have

$$(\bar{b}_3 \wedge 1_{K' \wedge M})f'_5 = 0. \quad (3.30)$$

By virtue of (3.2), we have

$$f'_5 = (\bar{a}_3 \wedge 1_{K' \wedge M})f'_6 \quad (3.31)$$

with $f'_6 \in [\Sigma^{p^n q + (p+2)q+2} M, E_4 \wedge K' \wedge M]$. By (3.27) and (3.2), we have

$$\begin{aligned} & (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r})d(f_2 i j) \\ &= (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M u_6) f'_3 \\ &= (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge u_6) f'_3, \quad \text{by (3.28)} \\ &= (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge j') f'_4, \quad \text{by (3.29)} \\ &= (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge j')(1_{E_3} \wedge \pi) f'_5, \quad \text{by (3.31)} \\ &= (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge j')(1_{E_3} \wedge \pi)(\bar{a}_3 \wedge 1_{K' \wedge M}) f'_6 \\ &= (\bar{a}_2 \bar{a}_3 \wedge 1_{K' \wedge M})(1_{E_4} \wedge (v \wedge 1_M) \bar{m}_M j')(1_{E_4} \wedge \pi) f'_6. \end{aligned}$$

That is,

$$(\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r})d(f_2 i j) = (\bar{a}_2 \bar{a}_3 \wedge 1_{K' \wedge M})(1_{E_4} \wedge (v \wedge 1_M) \bar{m}_M j')(1_{E_4} \wedge \pi) f'_6. \quad (3.32)$$

Since $[(\bar{b}_4 \wedge 1_K)(1_{E_4} \wedge \pi) f'_6] \in Ext_A^{4, p^n q + (p+2)q+2}(H^* K, H^* M) = 0$ (cf. [7, Proposition 2.2]), then by (3.1), we know that the d_1 -cycle $(\bar{b}_4 \wedge 1_K)(1_{E_4} \wedge \pi) f'_6$ is a d_1 -boundary. It follows that $(\bar{b}_4 \wedge 1_K)(1_{E_4} \wedge \pi) f'_6 = (\bar{b}_4 \wedge 1_K)(\bar{c}_3 \wedge 1_K) f'_7$ for some $f'_7 \in [\Sigma^{p^n q + (p+2)q+2} M, KG_3 \wedge K]$. Thus we have

$$(1_{E_4} \wedge \pi) f'_6 = (\bar{c}_3 \wedge 1_K) f'_7 + (\bar{a}_4 \wedge 1_K) f'_8 \quad (3.33)$$

with $f'_8 \in [\Sigma^{p^n q + (p+2)q+3} M, E_5 \wedge K]$. Then by (3.32), (3.33) and (3.2), we have

$$(\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r})d(f_2 i j) = (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge (v \wedge 1_M) \bar{m}_M j') f'_8. \quad (3.34)$$

Moreover, by composing $(1_{E_2} \wedge \bar{r})$ on (3.13) it is easy to get that

$$(\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r})d(f_2 i j) = (1_{E_2} \wedge \bar{r}(i'' \wedge 1_K) \beta) d(\eta'_{n,2} i' i j). \quad (3.35)$$

Combining (3.34) and (3.35) yields

$$(1_{E_2} \wedge \bar{r}(i'' \wedge 1_K) \beta) d(\eta'_{n,2} i' i j) = (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge (v \wedge 1_M) \bar{m}_M j') f'_8. \quad (3.36)$$

Notice that $\bar{r}(i'' \wedge 1_K) = (v \wedge 1_M) \bar{m}_M j'$ (see Lemma 3.3). Then (3.36) can turn into

$$(1_{E_2} \wedge (v \wedge 1_M) \bar{m}_M j' \beta) d(\eta'_{n,2} i' i j) = (1_{E_2} \wedge (v \wedge 1_M) \bar{m}_M j') (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_K) f'_8. \quad (3.37)$$

By (3.37) and (3.6), we have

$$(1_{E_2} \wedge j' \beta) d(\eta'_{n,2} i' i j) = (1_{E_2} \wedge j') (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_K) f'_8 + (1_{E_2} \wedge j' \alpha') f'_9 \quad (3.38)$$

with $f'_9 \in [\Sigma^{p^n q + (p+1)q+1} M, E_2 \wedge K]$. From [7, p.489], we know that the left hand side of (3.38) has filtration 4. However, since the first term of the right hand side of (3.38) has filtration ≥ 5 , then the second term of (3.38) must be of filtration 4. So f'_9 has filtration ≤ 3 . Notice the facts that $Ext_A^{3,p^n q + (p+1)q+2}(H^* K, H^* M) = 0$ (cf. Proposition 2.5) and $Ext_A^{2,p^n q + (p+1)q+1}(H^* K, H^* M) \cong \mathbb{Z}_p \{\beta_* i'_*(\tilde{h}_n)\}$ (cf. Proposition 2.7), then we have $(\bar{b}_2 \wedge 1_K) f'_9 = (1_{KG_2} \wedge \beta)(1_{KG_2} \wedge i')(\tilde{h}_n)$. Let $\xi_n = (\bar{a}_0 \bar{a}_1 \wedge 1_K) f'_9$. Then ξ_n is represented by $\beta_* i'_*(\tilde{h}_n)$ in the Adams spectral sequence. And so $\zeta_n = \xi_n i$ is represented by $i^* \beta_* i'_*(\tilde{h}_n) = \beta_* i'_* i^*(\tilde{h}_n) = \beta_* i'_* i_*(h_n) \neq 0 \in Ext_A^{2,p^n q + (p+1)q+1}(H^* K, \mathbb{Z}_p)$ (cf. Proposition 2.6). Thus Theorem I is proved.

Proof of Lemma 3.12. We first recall three cofibrations given in [7]

$$\Sigma^{-1} K \xrightarrow{vj'} \Sigma^q K' \xrightarrow{\bar{\psi}} K'_2 \xrightarrow{\bar{\rho}} K \quad (\text{see [7, (2.5)]}), \quad (3.39)$$

$$\Sigma^{q-1} K'_1 \xrightarrow{\varepsilon(1_{K'_1} \wedge i)} M \xrightarrow{w_2} X \xrightarrow{u_2} \Sigma^q K' \quad (\text{see [7, (3.7)]}), \quad (3.40)$$

$$X \xrightarrow{\bar{\psi} u_2} K'_2 \xrightarrow{w_3} K' \wedge W \xrightarrow{u_3} \Sigma X \quad (\text{see [7, (3.10)]}) \quad (3.41)$$

with the relation $u_2 u_3 = -v j' \pi$ [7, (3.11)]. By composing $(\bar{a}_2 \wedge 1_{K' \wedge M})$ on (3.27), we have

$$\begin{aligned} & (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r}) d(f_2 i j) \\ &= (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M u_6) f'_3, \quad \text{by (3.28)} \\ &= (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge j') f'_4, \quad \text{by (3.29)} \\ &= (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge j')(1_{E_3} \wedge \pi) f'_5. \end{aligned}$$

That is,

$$(\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r}) d(f_2 i j) = (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge j')(1_{E_3} \wedge \pi) f'_5. \quad (3.42)$$

By composing $(1_{E_2} \wedge (1_{K'} \wedge j))$ on (3.42), we have

$$\begin{aligned} & (1_{E_2} \wedge (1_{K'} \wedge j)) (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r}) d(f_2 i j) \\ &= (1_{E_2} \wedge (1_{K'} \wedge j)) (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge j')(1_{E_3} \wedge \pi) f'_5. \end{aligned} \quad (3.43)$$

On the one hand, for the left hand side of (3.43), we have

$$\begin{aligned} & (1_{E_2} \wedge (1_{K'} \wedge j)) (\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K' \wedge M})(1_{E_5} \wedge \bar{r}) d(f_2 i j) \\ &= -(\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K'}) (1_{E_5} \wedge (1_{K'} \wedge j) \bar{r}) d(f_2 i j), \quad \text{by Lemma 3.4} \\ &= -(\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K'}) (1_{E_5} \wedge \bar{\Delta}_{K'}) d(f_2 i j). \end{aligned}$$

On the other hand, for the right hand side of (3.43) we have

$$\begin{aligned} & (1_{E_2} \wedge (1_{K'} \wedge j)) (\bar{a}_2 \wedge 1_{K' \wedge M})(1_{E_3} \wedge (v \wedge 1_M) \bar{m}_M)(1_{E_3} \wedge j')(1_{E_3} \wedge \pi) f'_5 \\ &= -(\bar{a}_2 \wedge 1_{K'}) (1_{E_3} \wedge (1_{K'} \wedge j) (v \wedge 1_M) \bar{m}_M j' \pi) f'_5 \\ &= -(\bar{a}_2 \wedge 1_{K'}) (1_{E_3} \wedge (v \wedge 1_{s^0}) (1_M \wedge j) \bar{m}_M j' \pi) f'_5, \quad \text{since } (1_M \wedge j) \bar{m}_M = 1_M \\ &= -(\bar{a}_2 \wedge 1_{K'}) (1_{E_3} \wedge v j' \pi) f'_5, \quad \text{since } u_2 u_3 = -v j' \pi \\ &= (\bar{a}_2 \wedge 1_{K'}) (1_{E_3} \wedge u_2 u_3) f'_5. \end{aligned}$$

Thus we have

$$(\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K'}) (1_{E_5} \wedge \bar{\Delta}_{K'}) d(f_2 i j) = -(\bar{a}_2 \wedge 1_{K'}) (1_{E_3} \wedge u_2 u_3) f'_5. \quad (3.44)$$

Let X be the cofibre of $\varepsilon(1_{K'} \wedge i) : \Sigma^{q-1} K' \longrightarrow M$ given by the cofibration

$$\Sigma^{q-1} K' \xrightarrow{\varepsilon(1_{K'} \wedge i)} M \xrightarrow{w_2} X \xrightarrow{u_2} \Sigma^q K' \quad (\text{see [7, (3.7)]}).$$

[7, (3.8)] follows from [7, (3.7)] and [7, (3.6)] that

$$(\bar{a}_4 \wedge 1_{K'}) (1_{E_5} \wedge \bar{\Delta}_{K'}) d(f_2 i j) = (1_{E_4} \wedge u_2) f_3$$

for some $f_3 \in [\Sigma^{p^n q + (p+2)q+1} M, E_4 \wedge X]$. By composing $(\bar{a}_2 \bar{a}_3 \wedge 1_{K'})$ on [7, (3.8)], we have

$$(\bar{a}_2 \bar{a}_3 \bar{a}_4 \wedge 1_{K'}) (1_{E_5} \wedge \bar{\Delta}_{K'}) d(f_2 i j) = (\bar{a}_2 \bar{a}_3 \wedge 1_{K'}) (1_{E_4} \wedge u_2) f_3. \quad (3.45)$$

Combining (3.44) and (3.45) yields

$$(\bar{a}_2 \wedge 1_{K'}) (1_{E_3} \wedge u_2 u_3) f'_5 = -(\bar{a}_2 \bar{a}_3 \wedge 1_{K'}) (1_{E_4} \wedge u_2) f_3. \quad (3.46)$$

By [10, (1.7)], (3.46) can turn into

$$(1_{E_2} \wedge u_2) (\bar{a}_2 \wedge 1_X) (1_{E_3} \wedge u_3) f'_5 = -(1_{E_2} \wedge u_2) (\bar{a}_2 \bar{a}_3 \wedge 1_X) f_3. \quad (3.47)$$

From (3.47) and (3.40) we have

$$(\bar{a}_2 \wedge 1_X) (1_{E_3} \wedge u_3) f'_5 = -(\bar{a}_2 \bar{a}_3 \wedge 1_X) f_3 + (1_{E_2} \wedge w_2) \bar{f}_4 \quad (3.48)$$

with $\bar{f}_4 \in [\Sigma^{p^n q + (p+2)q-1} M, E_2 \wedge M]$.

Note that $(\bar{b}_2 \wedge 1_M) \bar{f}_4 \in [\Sigma^{p^n q + (p+2)q-1} M, KG_2 \wedge M] = 0$ by the exact sequence $[\Sigma^{p^n q + (p+2)q-1} M, KG_2] \xrightarrow{(1 \wedge i)^*} [\Sigma^{p^n q + (p+2)q-1} M, KG_2 \wedge M] \xrightarrow{(1 \wedge j)^*} [\Sigma^{p^n q + (p+2)q-2} M, KG_2]$ induced by (1.1), where the first and the last group are zero by the fact that $\pi_{p^n q + (p+2)q+r} KG_2 \cong Ext_A^{2, p^n q + (p+2)q+r}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $r = 0, -1, -2$ (cf. [2]). Hence, $\bar{f}_4 = (\bar{a}_2 \wedge 1_M) \bar{f}_5$ for some $\bar{f}_5 \in [\Sigma^{p^n q + (p+2)q} M, E_3 \wedge M]$. By (3.2) and (3.48), we have

$$(1_{E_3} \wedge u_3) f'_5 = -(\bar{a}_3 \wedge 1_X) f_3 + (1_{E_3} \wedge w_2) \bar{f}'_5 + (\bar{c}_2 \wedge 1_X) g_6 \quad (3.49)$$

with $g_6 \in [\Sigma^{p^n q + (p+2)q} M, KG_2 \wedge X]$. And so we have

$$(\bar{b}_3 \wedge 1_X) (1_{E_3} \wedge u_3) f'_5 = (\bar{b}_3 \wedge 1_X) (1_{E_3} \wedge w_2) \bar{f}'_5 + (\bar{b}_3 \bar{c}_2 \wedge 1_X) g_6. \quad (3.50)$$

Since $Ext_A^{3, p^n q + (p+2)q}(H^* M, H^* M) \cong \mathbb{Z}_p \{i_* j_* \overline{\bar{h}_n g_0}, j^* i^* \overline{\bar{h}_n g_0}\}$ (cf. Proposition 2.3), then we can have $(\bar{b}_3 \wedge 1_M) \bar{f}'_5 = \lambda_1 \overline{\bar{h}_n g_0} i j + \lambda_2 (1_{KG_3} \wedge i j) \overline{\bar{h}_n g_0}$ for some $\lambda_1, \lambda_2 \in \mathbb{Z}_p$, where $\overline{\bar{h}_n g_0} \in [\Sigma^{p^n q + (p+2)q+1} M, KG_3 \wedge M]$. And so

$$0 = \lambda_1 (\bar{c}_3 \wedge 1_M) \overline{\bar{h}_n g_0} i j + \lambda_2 (\bar{c}_3 \wedge 1_M) (1_{KG_3} \wedge i j) \overline{\bar{h}_n g_0}.$$

By composing i on the above equality, we get that $\lambda_2(\bar{c}_3 \wedge 1_M)(1_{KG_3} \wedge ij)\overline{\overline{h_n g_0}} i = 0$. However, since $d_2(i^*(ij)_*\overline{\overline{h_n g_0}}) = i^*d_2(i_*j_*\overline{\overline{h_n g_0}}) \neq 0$ (see Proposition 2.4 (1)), we get that $(\bar{c}_3 \wedge 1_M)(1_{KG_3} \wedge ij)\overline{\overline{h_n g_0}} i \neq 0$. Thus, we have that

$$\lambda_2 = 0, \lambda_1(\bar{c}_3 \wedge 1_M)\overline{\overline{h_n g_0}} ij = 0.$$

Note that $d_2(j^*i^*\overline{\overline{h_n g_0}}) \neq 0$ by Proposition 2.4 (2), then $(\bar{c}_3 \wedge 1_M)\overline{\overline{h_n g_0}} ij \neq 0$. Thus we see that $\lambda_1 = 0$. From the above discussion, we know that $(\bar{b}_3 \wedge 1_M)\bar{f}_5 = 0$ and (3.50) can turn into

$$(\bar{b}_3 \wedge 1_X)(1_{E_3} \wedge u_3)f'_5 = (\bar{b}_3 \bar{c}_2 \wedge 1_X)g_6. \quad (3.51)$$

The argument of the proof from [7, (3.16)] to [7, p.491] shows that $(\bar{b}_3 \wedge 1_X)(1_{E_3} \wedge u_3)f_6 = -(\bar{b}_3 \bar{c}_2 \wedge 1_X)\tilde{l}_0$ in [7, (3.16)] implies $(\bar{b}_3 \wedge 1_{K' \wedge M})f_6 = 0$. By a similar argument as in [7], we can also show that (3.51) implies (3.30) holds. $\square$

Proof of Theorem II By Theorem I, we get that

$$\beta_* i'_* i_*(h_n) \neq 0 \in Ext_A^{2, p^n q + (p+1)q+1}(H^*K, \mathbb{Z}_p)$$

is a permanent cycle in the Adams spectral sequence and converges to a nontrivial element $\zeta_n \in \pi_{p^n q + (p+1)q-1}K$.

Consider the following composition of maps

$$\bar{f} : \Sigma^{p^n q + (p+1)q-1} S \xrightarrow{\zeta_n} K \xrightarrow{jj'\beta} \Sigma^{-pq+2} S.$$

Since ζ_n is represented up to nonzero scalar by $\beta_* i'_* i_*(h_n) \in Ext_A^{2, p^n q + (p+1)q+1}(H^*K, \mathbb{Z}_p)$ in the Adams spectral sequence, then the above $\bar{f}$ is represented up to nonzero scalar by $\bar{c} = (jj'\beta)_* \beta_* i'_* i_*(h_n)$ in the Adams spectral sequence.

Meanwhile, it is well known that the β -element $\beta_2 = jj'\beta^2 i'_* i$ is represented by $k_0 \in Ext_A^{2, 2pq+q}(\mathbb{Z}_p, \mathbb{Z}_p)$ in the Adams spectral sequence. By the knowledge of Yoneda products we can see that $\bar{f}$ is represented (up to nonzero scalar) by

$$\bar{c} = k_0 h_n \neq 0 \in Ext_A^{3, q(p^n+2p+1)}(\mathbb{Z}_p, \mathbb{Z}_p)$$

in the Adams spectral sequence(cf. [6, Table 8.1]).

Moreover, we know that $Ext_A^{3-r, q(p^n+2p+1)+(-r+1)}(\mathbb{Z}_p, \mathbb{Z}_p) = 0$ for $r \geq 2$, then $k_0 h_n$ cannot be hit by any differential in the Adams spectral sequence, and so the corresponding homotopy element $\bar{f} \in \pi_* S$ is nontrivial and of order p . This finishes the proof of Theorem II. $\square$

References

1. Cohen R. Odd primary families in stable homotopy theory. *Memoirs of the American Mathematical Society*, 1981, **242**.

2. Liulevicius A. The factorizations of cyclic reduced powers by secondary cohomology operations. *Memoirs of the American Mathematical Society*, 1962, **42**.
3. Wang X J. Ordered cochain complex and $H^{4,*}(A_p)$ for $p \geq 5$. *Beijing Mathematics*, 1995, 1: 80 ~ 99.
4. Liu X G. A nontrivial product in the stable homotopy groups of spheres. *Sci. in China Ser. A*, 2004, **47**(6): 831 ~ 841.
5. H. Toda. On spectra realizing exterior part of the Steenrod algebra. *Topology*, 1971, **10**: 53 ~ 65.
6. Aikawa T. 3-dimensional cohomology of the mod p Steenrod algebra. *Mathematica Scandinavica*, 1980, **47**: 91 ~ 115.
7. Lin J K. A new family of filtration three in the stable homotopy of spheres. *Hiroshima Mathematical Journal*, 2001, **31**: 477 ~ 492.
8. Lin J K and Zheng Q B. A new family of filtration seven in the stable homotopy of spheres. *Hiroshima Mathematical Journal*, 1998, **28**: 183 ~ 205.
9. Cohen R. and Goerss P. Secondary cohomology operations that detect homotopy classes, *Topology*, 1984, **23**: 177 – 194.
10. Toda H. Algebra of stable homotopy of $\mathbb{Z}_p$ -spaces and applications. *Journal of Mathematics of Kyoto University*, 1971, **11**: 197 ~ 251.
11. Oka S. Multilicative structure of finite ring spectra and stable homotopy of spheres. *Algebraic Topology (Aarhus), Lecture Notes in Mathematics*, 1984, **1051**: 418 ~ 441. Springer-Verlag.
12. Thomas E. and Zahler R. Generalized higher order cohomology operations and stable homotopy groups of spheres. *Advances in Mathematics*, 1976, **20**: 287 ~ 328.